Applicant: Frederick L. Hall, et al. Attorney's Docket No.: 14230-010002 / 2895

Serial No. : 10/733,852

Filed December 10, 2003

Page 2 of 8

## AMENDMENTS TO THE CLAIMS

This listing of claims replaces all prior versions and listings of claims in the application:

## Listing of Claims:

- 1.-65. (Canceled)
- 66. (Currently Amended) A fusion polypeptide comprising a collagen-binding domain and an epithelial cell proliferation-modulating agent, wherein the epithelial cell proliferation-modulating agent is selected from the group consisting of hemopoietic growth factors (HeGFs), insulin, nerve growth factor (NGF), NGF receptor, epidermal growth factor (EGF) receptor, neu, inhibin α, inhibin β, Müllerian inhibitory substance, tumor necrosis factor (TNF)-receptor (type 1), TNF-receptor (type 2), platelet-derived growth factor (PDGF) receptor α, PDGF receptor β, wnt-2, hst/ks3, and hepatocyte growth factor (HGF) receptor (c-met), stem-cell-factor (SCF), SCF receptor (c-kit), erythropoietin (epo), epo-receptor, and leukemia inhibitory factor.
- 67. (Previously presented) The fusion polypeptide of claim 66, wherein the epithelial cell proliferation-modulating agent stimulates epithelial cell proliferation.
- 68. (Previously presented) The fusion polypeptide of Claim 66, wherein the collagenbinding domain is a collagen-binding domain of von Willebrand factor.
- 69. (Previously presented) The fusion polypeptide of claim 68, wherein the collagen-binding domain of von Willebrand factor comprises the decapeptide WREPSFMALS (SEQ ID NO:1).
- 70. (Canceled)
- 71. (Canceled)

Applicant: Frederick L. Hall, et al. Attorney's Docket No.: 14230-010002 / 2895

Serial No. : 10/733,852

Filed December 10, 2003

Page : 3 of 8

72. (Currently Amended) A nucleic acid sequence encoding a fusion polypeptide comprising a collagen-binding domain and an epithelial cell proliferation-modulating agent, wherein the epithelial cell proliferation-modulating agent is selected from the group consisting of hemopoietic growth factors (HeGFs), insulin, nerve growth factor (NGF), NGF receptor, epidermal growth factor (EGF) receptor, neu, inhibin α, inhibin β, Müllerian inhibitory substance, tumor necrosis factor (TNF)-receptor (type 1), TNF-receptor (type 2), platelet-derived growth factor (PDGF) receptor α, PDGF receptor β, wnt-2, hst/ks3, and hepatocyte growth factor (HGF) receptor (c-met), stem cell-factor (SCF), SCF receptor (c-kit), erythropoietin (epo), epo receptor, and leukemia inhibitory factor.

- 73. (Previously presented) The nucleic acid sequence of claim 72, operably linked to a promoter.
- 74. (Previously presented) An expression vector comprising the nucleic acid sequence of claim 72.
- 75. (Previously presented) The expression vector of claim 74, wherein the expression vector is a retroviral vector.
- 76. (Previously presented) A host cell comprising the nucleic acid sequence of claim 72.
- 77. (Previously presented) A method of producing the fusion polypeptide comprising a collagen-binding domain and an epithelial cell proliferation-modulating agent, comprising growing the host cells of claim 76 under conditions that allow expression of the fusion polypeptide recovering the fusion polypeptide.
- 78. (Previously presented) The method of claim 77, wherein the host is a prokaryotic cell.
- 79. (Previously presented) The method of claim 77, wherein the host is a eukaryotic cell.

Applicant : Frederick L. Hall, et al. Attorney's Docket No.: 14230-010002 / 2895

Serial No. : 10/733,852

Filed : December 10, 2003

Page : 4 of 8

80. (Currently Amended) A pharmaceutical composition comprising a fusion polypeptide comprising a collagen-binding domain and an epithelial cell proliferation-modulating agent, in a pharmaceutically acceptable carrier, wherein the epithelial cell proliferation-modulating agent is selected from the group consisting of hemopoietic growth factors (HeGFs), insulin, nerve growth factor (NGF), NGF receptor, epidermal growth factor (EGF) receptor, neu, inhibin α, inhibin β, Müllerian inhibitory substance, tumor necrosis factor (TNF)-receptor (type 1), TNF-receptor (type 2), platelet-derived growth-factor (PDGF) receptor α, PDGF receptor β, wnt-2, hst/ks3, and hepatocyte growth factor (HGF) receptor (c-met), stem cell-factor (SCF), SCF receptor (c-kit), erythropoietin (epo), epo-receptor, and leukemia inhibitory factor.